

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	4	(IGFBP adj "4") near4 (protease or peptidase or proteinase)	USPAT	OR	OFF	2005/11/09 18:59
L2	0	L1 near10 (antibody or antibodies)	USPAT	OR	OFF	2005/11/09 18:58
L3	0	L1 near10 (antibody or antibodies or monoclonal or polyclonal)	USPAT	OR	OFF	2005/11/09 18:59

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623SQS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *
* *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JUL 20 Powerful new interactive analysis and visualization
software,

STN AnaVist, now available
NEWS 4 AUG 11 STN AnaVist workshops to be held in North America
NEWS 5 AUG 30 CA/CAPlus -Increased access to 19th century
research documents

NEWS 6 AUG 30 CASREACT - Enhanced with displayable reaction
conditions

NEWS 7 SEP 09 ACD predicted properties enhanced in
REGISTRY/ZREGISTRY

NEWS 8 OCT 03 MATHDI removed from STN

NEWS 9 OCT 04 CA/CAPlus-Canadian Intellectual Property Office
(CIPO) added

to core patent offices
NEWS 10 OCT 06 STN AnaVist workshops to be held in North America
NEWS 11 OCT 13 New CAS Information Use Policies Effective October
17, 2005

NEWS 12 OCT 17 STN(R) AnaVist(TM), Version 1.01, allows the
export/download of CAPlus documents for use in third-party analysis
and

visualization tools
NEWS 13 OCT 27 Free KWIC format extended in full-text databases
NEWS 14 OCT 27 DIOGENES content streamlined
NEWS 15 OCT 27 EPFULL enhanced with additional content

NEWS EXPRESS JUNE 13 CURRENT WINDOWS VERSION IS V8.0, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 13 JUNE 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *
* *

FILE 'HOME' ENTERED AT 19:00:34 ON 09 NOV 2005

=> File Medline EMBASE Biosis Caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 19:00:41 ON 09 NOV 2005

FILE 'EMBASE' ENTERED AT 19:00:41 ON 09 NOV 2005
Copyright (c) 2005 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 19:00:41 ON 09 NOV 2005
Copyright (c) 2005 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 19:00:41 ON 09 NOV 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (IGFBP-4) (4A) (protease or peptidase or proteinase)
L1 358 (IGFBP-4) (4A) (PROTEASE OR PEPTIDASE OR PROTEINASE)

=> s L1 (10A) (antibody or antibodies or monoclonal or polyclonal)
L2 27 L1 (10A) (ANTIBODY OR ANTIBODIES OR MONOCLONAL OR
POLYCLONAL)

=> duplicate
ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove
ENTER L# LIST OR (END):l2
DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, BIOSIS, CAPLUS'
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n
PROCESSING COMPLETED FOR L2
L3 8 DUPLICATE REMOVE L2 (19 DUPLICATES REMOVED)

=> d l3 1-8 bib ab

L3 ANSWER 1 OF 8 MEDLINE on STN DUPLICATE 1
 AN 2004362217 MEDLINE
 DN PubMed ID: 15087430
 TI Pregnancy-associated plasma protein-a production in rat
 granulosa cells:
 stimulation by follicle-stimulating hormone and inhibition by the
 oocyte-derived bone morphogenetic protein-15.
 AU Matsui Motozumi; Sonntag Barbara; Hwang Seong Soo; Byerly Tara;
 Hourvitz
 Ariel; Adashi Eli Y; Shimasaki Shunichi; Erickson Gregory F
 CS Department of Reproductive Medicine, University of California
 San Diego,
 La Jolla, California 92093-0674, USA.
 NC U54 HD12303 (NICHD)
 SO Endocrinology, (2004 Aug) 145 (8) 3686-95. Electronic
 Publication:
 2004-04-15.
 Journal code: 0375040. ISSN: 0013-7227.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Abridged Index Medicus Journals; Priority Journals
 EM 200408
 ED Entered STN: 20040722
 Last Updated on STN: 20040818
 Entered Medline: 20040817
 AB Pregnancy-associated plasma protein-A (PAPP-A) is the major IGF
 binding
 protein-4 (IGFBP-4) protease in follicular fluid, consistent
 with its
 proposed role in folliculogenesis. Despite growing interest,
 almost
 nothing is known about how PAPP-A expression is regulated in any
 tissue.
 Here we show that FSH and oocytes regulate PAPP-A expression in
 granulosa
 cells (GCs). By in situ hybridization, ovary PAPP-A mRNA was
 markedly
 increased by pregnant mare serum gonadotropin treatment, and the
 message
 was localized to the membrana GCs but not cumulus GCs (CGCs) of
 dominant
 follicles. To explore the mechanism, we used primary cultures
 of rat GCs.
 Control (untreated) cells produced little or no PAPP-A
 spontaneously.
 Conversely, FSH markedly stimulated PAPP-A mRNA and protein in a
 dose- and
 time-dependent fashion. Interestingly, PAPP-A expression in
 isolated CGCs
 was also strongly induced by FSH, and the induction was
 inhibited by added

oocytes. To investigate the nature of the inhibition, we tested the effect of oocyte-derived bone morphogenetic protein-15 (BMP-15). BMP-15 alone had no effect on basal levels of PAPP-A expression by cultures of membrana GCs or CGCs. However, BMP-15 markedly inhibited the FSH stimulation of PAPP-A production in a dose-dependent manner. The cleavage of IGFBP-4 by conditioned media from FSH-treated GCs was completely inhibited by anti-PAPP-A **antibody**, indicating the **IGFBP -4 protease** secreted by GCs is PAPP-A. These results demonstrate stimulatory and inhibitory roles for FSH and BMP-15, respectively, in regulating PAPP-A production by GCs. We propose that FSH and oocyte-derived BMP-15 form a controlling network that ensures the spatiotemporal pattern of GC PAPP-A expression in the dominant follicle.

L3 ANSWER 2 OF 8 MEDLINE on STN DUPLICATE 2
 AN 2002654650 MEDLINE
 DN PubMed ID: 12414897
 TI Pregnancy-associated plasma protein A proteolytic activity is associated with the human placental trophoblast cell membrane.
 AU Sun Irene Y C; Overgaard Michael T; Oxvig Claus; Giudice Linda C
 CS Department of Gynecology and Obstetrics, Center for Research on Women's Health and Reproductive Medicine, Stanford University Medical Center,
 Stanford, CA 94305-5317, USA.
 NC HD 35789-05 (NICHD)
 SO Journal of clinical endocrinology and metabolism, (2002 Nov) 87 (11) 5235-40.
 Journal code: 0375362. ISSN: 0021-972X.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Abridged Index Medicus Journals; Priority Journals
 EM 200212
 ED Entered STN: 20021105
 Last Updated on STN: 20021217
 Entered Medline: 20021209
 AB Pregnancy-associated plasma protein-A (PAPP-A) is a product of the placenta and decidua and is secreted into the maternal circulation during human pregnancy. It recently has been identified as an IGF binding

protein (IGFBP)-4 protease. Presumed functions at the maternal-fetal interface are to proteolyze IGFBP-4 and thus increase IGF bioavailability locally in the placenta, to promote IGF-II-mediated trophoblast invasion into the maternal decidua, and to modulate IGF regulation of steroidogenesis and glucose and amino acid transport in the villous.

Herein, we have investigated the possibility that IGFBP-4 proteolysis may occur on the trophoblast cell membrane, presumably to increase local bioavailable IGF for interactions with cognate IGF membrane receptors.

Human trophoblasts were cultured, trophoblast plasma membranes were isolated and solubilized, and IGFBP-4 protease activity and PAPP-A immunoreactivity in the solubilized plasma membrane fraction were investigated. IGFBP-4 protease activity was detected in solubilized human trophoblast membranes, resulting in cleavage of recombinant human IGFBP-4 into 18- and 14-kDa fragments, detected by Western immunoblot analysis.

This protease activity was dependent on the presence of IGF-II, and its metal ion dependence was demonstrated by inhibition of the protease by the metal chelators, EDTA and EGTA. The presence of PAPP-A in solubilized human trophoblast membranes was demonstrated by Western immunoblotting.

Trophoblast membrane PAPP-A had a relative molecular weight of approximately 200 kDa and comigrated on (reducing) SDS-PAGE with recombinant human PAPP-A and PAPP-A secreted into media conditioned by cultured human trophoblasts. IGFBP-4 protease activity was not detected after immunodepletion of PAPP-A from the trophoblast membrane fraction with PAPP-A **polyclonal antibodies**, suggesting the identity of the membrane-derived **IGFBP-4 protease** as PAPP-A. Immunocytochemistry revealed PAPP-A on the cell membrane and in the cytoplasm of human trophoblasts in culture.

Together, these data demonstrate the presence of an IGF-II- and metal-dependent IGFBP-4 protease activity in human trophoblast plasma membranes, identified as PAPP-A, which is well situated to proteolyze

IGFBP-4 at the maternal-placental interface to facilitate IGF action at the villous surface and/or the invading extravillous cytotrophoblast.

L3 ANSWER 3 OF 8 MEDLINE on STN DUPLICATE 3
AN 2002255160 MEDLINE
DN PubMed ID: 11994388
TI Identification and regulation of the IGFBP-4 protease and its physiological inhibitor in human trophoblasts and endometrial stroma:
evidence for paracrine regulation of IGF-II bioavailability in the placental bed during human implantation.
AU Giudice L C; Conover C A; Bale L; Faessen G H; Ilg K; Sun I; Imani B; Suen L-F; Irwin J C; Christiansen M; Overgaard M T; Oxvig C
CS Department of Gynecology and Obstetrics, Stanford University Medical Center, Stanford, California 94305, USA.. giudice@stanford.edu
NC U54 HD31398 (NICHD)
SO Journal of clinical endocrinology and metabolism, (2002 May) 87 (5) 2359-66.
Journal code: 0375362. ISSN: 0021-972X.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals
EM 200206
ED Entered STN: 20020508
Last Updated on STN: 20020615
Entered Medline: 20020614
AB The IGF family plays an important role in implantation and placental physiology. IGF-II is abundantly expressed by placental trophoblasts, and IGF binding protein (IGFBP)-4, a potent inhibitor of IGF actions, is the second most abundant IGFBP in the placental bed, expressed exclusively by the maternal decidua. Proteolysis of IGFBP-4 results in decreased affinity for IGF peptides, thereby enhancing IGF actions. In the current study, we have identified the IGFBP-4 protease and its inhibitor in human trophoblast and decidualized endometrial stromal cell cultures, and we have investigated their regulation in an effort to understand control of IGF-II bioavailability at the placental-decidual interface in human

implantation. IGFBP-4 protease activity was detected in conditioned media

(CM) from human trophoblasts and decidualized endometrial stromal cells

using (125)I-IGFBP-4 substrate. Identification of the IGFBP-4 protease as

pregnancy-associated plasma protein-A (PAPP-A) was confirmed by specific

immunoinhibition and immunodepletion of the **IGFBP-4**

protease activity with specific PAPP-A **antibodies**. The

IGFBP-4 protease activity was IGF-II-dependent in trophoblast

CM. In

decidualized stromal CM, PAPP-A/IGFBP-4 protease activity was also

IGF-II-dependent, but was evident only when IGF-II was added in molar

excess of the predominant IGFBP in decidualized stromal cell CM, IGFBP-1,

supporting bioavailable IGF-II as a key cofactor of IGFBP-4 proteolysis by

PAPP-A. Cultured first and second trimester human trophoblasts (n = 5)

secreted PAPP-A into CM with mean +/- SEM levels of 172.4 +/- 32.8

mIU/liter.10(5) cells, determined by specific ELISA. PAPP-A in trophoblast CM (n = 3) and did not change in the presence of IGF-II (1-100

ng/ml). Cultured human endometrial stromal cells (n = 4) secreted low

levels of PAPP-A (6.25 +/- 3.6 mIU/liter.10(5) cells). A physiological

inhibitor of PAPP-A, the proform of eosinophil major basic protein

(promBP), was detected in trophoblast CM at levels of 1853 +/- 308

mIU/liter.10(5) cells, determined by specific ELISA, and was nearly

undetectable in CM of human endometrial stromal cells. Upon in vitro

decidualization of endometrial stromal cells with progesterone, PAPP-A

levels in CM increased nearly 9-fold without a concomitant change in

promBP. In contrast to the experiments with trophoblasts, IGF-II and the

IGF analogues, Leu(27) IGF-II, and Des (1-6) IGF-II, resulted in a

dose-dependent decrease of PAPP-A levels in decidualized endometrial

stromal CM by 70-90%, and a dose-dependent increase in promBP of 14- to

41-fold. The data demonstrate conclusively that the IGF-II-dependent

IGFBP-4 protease of human trophoblast and decidual origin is PAPP-A.

Furthermore, the differential regulation of decidual PAPP-A and proMBP by

insulin-like peptides supports a role for trophoblast-derived IGF-II as a

paracrine regulator of these maternal decidual products that have the

potential to regulate IGF-II bioavailability at the trophoblast-decidual

interface. Overall, the data underscore potential roles for a complex

family of enzyme (PAPP-A), substrate (IGFBP-4), inhibitor (proMBP), and

cofactor (IGF-II) in the placental bed during human implantation.

L3 ANSWER 4 OF 8 EMBASE COPYRIGHT (c) 2005 Elsevier B.V. All rights

reserved on STN

DUPLICATE 4

AN 2001166450 EMBASE

TI Pregnancy-associated plasma protein-A is the insulin-like growth factor

binding protein-4 protease secreted by human ovarian granulosa cells and

is a marker of dominant follicle selection and the corpus luteum.

AU Conover C.A.; Faessen G.F.; Ilg K.E.; Chandrasekher Y.A.; Christiansen M.;

Overgaard M.T.; Oxvig C.; Giudice L.C.

CS Dr. L.C. Giudice, Dept GYN, Stanford University Medical Center, Stanford,

CA 94305, United States

SO Endocrinology, (2001) Vol. 142, No. 5, pp. 2155-2158.

Refs: 24

ISSN: 0013-7227 CODEN: ENDOAO

CY United States

DT Journal; Article

FS 003 Endocrinology

010 Obstetrics and Gynecology

029 Clinical Biochemistry

LA English

SL English

ED Entered STN: 20010523

Last Updated on STN: 20010523

AB Insulin-like growth factors (IGFs), IGF binding proteins (IGFBPs), and

IGFBP proteases are important in ovarian function. IGFs stimulate

granulosa steroidogenesis, an effect that is inhibited by IGFBP-4 and

augmented by IGFBP-4 proteolysis. We have recently identified the IGFBP-4

protease in human ovarian follicular fluid (FF) as pregnancy-associated plasma protein-A (PAPP-A). In the current study, we identify the IGFBP-4 protease secreted by cultured human ovarian granulosa cells as PAPP-A, based on specific immunoinhibition and immunodepletion of the IGFBP-4 protease activity with PAPP-A polyclonal antibodies and immunorecognition by PAPP-A monoclonal antibodies in ELISA. PAPP-A was barely detectable in conditioned media (CM) from granulosa derived from ≤ 9 mm androgendominant follicles, but was secreted by cultured granulosa from estrogen-dominant follicles ≥ 9 mm, coincident with dominant follicle selection, and by luteinizing granulosa. PAPP-A levels in CM from the latter did not change in response to IGF-II or hCG (100 ng/mL). A naturally occurring inhibitor of PAPP-A, proform of eosinophil major basic protein (promBP), was detected by ELISA in estrogen-dominant follicular fluid FF, but not in CM from granulosa or luteinizing granulosa cells treated with IGF-II (0-200 ng/mL), FSH (0-100 ng/mL) or hCG (0-100 ng/mL), suggesting an alternative source (other than granulosa) for promBP, compared to PAPP-A. The data demonstrate granulosa cells as a source of PAPP-A in human ovary and suggest that PAPP-A is a marker of ovarian follicle selection and corpus luteum formation. In addition the data suggest complex regulation of this system in human ovary.

L3 ANSWER 5 OF 8 MEDLINE on STN

AN 2001262826 MEDLINE

DN PubMed ID: 11316785

TI Pregnancy-associated plasma protein-a is the insulin-like growth factor

binding protein-4 protease secreted by human ovarian granulosa cells and

is a marker of dominant follicle selection and the corpus luteum.

AU Conover C A; Faessen G F; Ilg K E; Chandrasekher Y A; Christiansen M;

Overgaard M T; Oxvig C; Giudice L C

CS Endocrine Unit, Mayo Clinic, Rochester, Minnesota 55905, USA.

NC HD31579-07 (NICHD)

SO Endocrinology, (2001 May) 142 (5) 2155.

Journal code: 0375040. ISSN: 0013-7227.

CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Abridged Index Medicus Journals; Priority Journals
 EM 200105
 ED Entered STN: 20010521
 Last Updated on STN: 20010521
 Entered Medline: 20010517
 AB Insulin-like growth factors (IGFs), IGF binding proteins (IGFBPs), and IGFBP proteases are important in ovarian function. IGFs stimulate granulosa steroidogenesis, an effect that is inhibited by IGFBP-4 and augmented by IGFBP-4 proteolysis. We have recently identified the IGFBP-4 protease in human ovarian follicular fluid (FF) as pregnancy-associated plasma protein-A (PAPP-A). In the current study, we identify the IGFBP-4 protease secreted by cultured human ovarian granulosa cells as PAPP-A, based on specific immunoinhibition and immunodepletion of the **IGFBP-4 protease** activity with PAPP-A **polyclonal antibodies** and immunorecognition by PAPP-A monoclonal antibodies in ELISA. PAPP-A was barely detectable in conditioned media (CM) from granulosa derived from ≤ 9 mm androgen-dominant follicles, but was secreted by cultured granulosa from estrogen-dominant follicles ≥ 9 mm, coincident with dominant follicle selection, and by luteinizing granulosa. PAPP-A levels in CM from the latter did not change in response to IGF-II or hCG (100 ng/mL).
 A naturally occurring inhibitor of PAPP-A, proform of eosinophil major basic protein (promBP), was detected by ELISA in estrogen-dominant follicular fluid FF, but not in CM from granulosa or luteinizing granulosa cells treated with IGF-II (0-200 ng/mL), FSH (0-100 ng/mL) or hCG (0-100 ng/mL), suggesting an alternative source (other than granulosa) for promBP, compared to PAPP-A. The data demonstrate granulosa cells as a source of PAPP-A in human ovary and suggest that PAPP-A is a marker of ovarian follicle selection and corpus luteum formation. In addition the data suggest complex regulation of this system in human ovary.

L3 ANSWER 6 OF 8 MEDLINE on STN DUPLICATE 5
 AN 2001260195 MEDLINE
 DN PubMed ID: 11231911
 TI Insulin-like growth factor binding protein-4 protease produced
 by smooth muscle cells increases in the coronary artery after angioplasty.
 AU Bayes-Genis A; Schwartz R S; Lewis D A; Overgaard M T;
 Christiansen M;
 Oxvig C; Ashai K; Holmes D R Jr; Conover C A
 CS Division of Cardiovascular Diseases, Mayo Clinic and Foundation,
 Rochester, Minnesota, USA.
 SO Arteriosclerosis, thrombosis, and vascular biology, (2001 Mar)
 21 (3)
 335-41.
 Journal code: 9505803. ISSN: 1524-4636.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200105
 ED Entered STN: 20010521
 Last Updated on STN: 20010521
 Entered Medline: 20010517
 AB Insulin-like growth factor (IGF)-I stimulates vascular smooth
 muscle cell
 (VSMC) migration and proliferation, which are fundamental to
 neointimal
 hyperplasia in postangioplasty restenosis. IGF-I action is
 modulated by
 several high-affinity IGF binding proteins (IGFBPs). IGFBP-4 is
 the
 predominant IGFBP produced by VSMCs and is a potent inhibitor of
 IGF-I
 action. However, specific IGFBP-4 proteases can cleave IGFBP-4
 and
 liberate active IGF-I. In this study, we document IGFBP-4
 protease
 produced by human and porcine coronary artery VSMCs in culture as
 pregnancy-associated plasma protein-A (PAPP-A). This was shown
 by a
 distinctive IGFBP-4 cleavage pattern, specific inhibition of
 IGFBP
 -4 protease activity with PAPP-A polyclonal
 antibodies, and immunorecognition of PAPP-A by monoclonal
 antibodies. Furthermore, we found a 2-fold increase in IGFBP-4
 protease
 activity in injured porcine VSMC cultures in vitro ($P < 0.05$). We
 also
 evaluated IGFBP-4 protease/PAPP-A expression in vivo after
 coronary artery
 balloon injury. Twenty-five immature female pigs underwent
 coronary

overstretch balloon injury, and vessels were examined at defined time points after the procedure. Abundant PAPP-A expression was observed in the cytoplasm of medial and neointimal cells 7, 14, and 28 days after angioplasty ($P < 0.01$ vs control). The highest PAPP-A labeling indices were located in the neointima ($36.1 \pm 2.1\%$) and the media ($31.7 \pm 1.2\%$) 28 days after injury. Western blot analysis confirmed increased PAPP-A in injured vessels. PAPP-A, a regulator of IGF-I bioavailability through proteolysis of IGFBP-4, is thus expressed by VSMCs in vitro and in restenotic lesions in vivo. These results suggest a possible role for PAPP-A in neointimal hyperplasia.

L3 ANSWER 7 OF 8 MEDLINE on STN DUPLICATE 6
AN 2000066728 MEDLINE
DN PubMed ID: 10599745
TI Evidence that the insulin-like growth factor binding protein-4 protease in human ovarian follicular fluid is pregnancy associated plasma protein-A.
AU Conover C A; Oxvig C; Overgaard M T; Christiansen M; Giudice L C
CS Endocrine Research Unit, Mayo Clinic and Mayo Foundation, Rochester, MN 55905, USA.. Conover.Cheryl@Mayo.Edu
NC HD 31579-05 (NICHD)
SO Journal of clinical endocrinology and metabolism, (1999 Dec) 84 (12) 4742-5.
Journal code: 0375362. ISSN: 0021-972X.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals
EM 200001
ED Entered STN: 20000124
Last Updated on STN: 20000124
Entered Medline: 20000107
AB Ovarian insulin-like growth factor binding protein-4 (IGFBP-4) proteolysis is involved in the regulation of follicular development, but until now the identity of the responsible enzyme was unknown. In this study, we identify the IGFBP4 protease in human follicular fluid as pregnancy

associated plasma protein-A (PAPP-A) based on distinctive IGFBP-4 cleavage pattern, the same protease inhibitor profile, specific inhibition and immunodepletion of **IGFBP-4 protease** activity with PAPP-A **polyclonal antibodies**, and immunorecognition by PAPP-A monoclonal antibodies in ELISA. Furthermore, PAPP-A levels in estrogen-dominant and androgen-dominant follicular fluids reflect their IGFBP-4 proteolytic activity. PAPP-A was also secreted by human granulosa cells, the reputed source of IGFBP-4 protease activity in follicular fluid. We have the molecular and biochemical tools to begin to delineate the regulation and biological function of PAPP-A in normal and dysregulated follicular development and atresia.

L3 ANSWER 8 OF 8 MEDLINE on STN DUPLICATE 7
 AN 1999179030 MEDLINE
 DN PubMed ID: 10077652
 TI The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A.
 AU Lawrence J B; Oxvig C; Overgaard M T; Sottrup-Jensen L; Gleich G J; Hays L G; Yates J R 3rd; Conover C A
 CS Endocrine Research Unit, Mayo Clinic and Mayo Foundation, Rochester, MN 55905, USA.
 NC AI09728 (NIAID)
 DK07352 (NIDDK)
 GM52095 (NIGMS)
 SO Proceedings of the National Academy of Sciences of the United States of America, (1999 Mar 16) 96 (6) 3149-53.
 Journal code: 7505876. ISSN: 0027-8424.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199905
 ED Entered STN: 19990601
 Last Updated on STN: 20000303
 Entered Medline: 19990520
 AB Proteolytic cleavage of the six known insulin-like growth factor binding proteins (IGFBPs) is a powerful means of rapid structure and function

modification of these important growth-regulatory proteins.

Intact

IGFBP-4 is a potent inhibitor of IGF action in vitro, and cleavage of

IGFBP-4 has been shown to abolish its ability to inhibit IGF stimulatory

effects in a variety of systems, suggesting that IGFBP-4 proteolysis acts

as a positive regulator of IGF bioavailability. Here we report the

isolation of an IGF-dependent IGFBP-4-specific protease from human

fibroblast-conditioned media and its identification by mass spectrometry

microsequencing as pregnancy-associated plasma protein-A (PAPP-A), a

protein of unknown function found in high concentrations in the maternal

circulation during pregnancy. **Antibodies** raised against PAPP-A both inhibited and immunodepleted **IGFBP-4**

protease activity in human fibroblast-conditioned media.

Moreover, PAPP-A purified from pregnancy sera had IGF-dependent IGFBP-4

protease activity. PAPP-A mRNA was expressed by the human fibroblasts and


osteoblasts, and PAPP-A protein was secreted into the culture medium. In

conclusion, we have identified an IGF-dependent IGFBP protease and at the

same time assigned a function to PAPP-A. This represents an unanticipated

union of two areas of research that were not linked in any way before this

report.



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Display Show Sort by Send to

About Entrez

All: 49 Review: 2

Text Version

Items 1 - 20 of 49

Page of 3 Next

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

☐ 1: [Jia D, Heersche JN.](#)

[Related Articles, Links](#)

Pregnancy-associated plasma protein-A proteolytic activity in rat vertebral cell cultures: stimulation by dexamethasone--a potential mechanism for glucocorticoid regulation of osteoprogenitor proliferation and differentiation.

J Cell Physiol. 2005 Sep;204(3):848-58.

PMID: 15754336 [PubMed - indexed for MEDLINE]

☐ 2: [Qin QP, Kokkala S, Lund J, Tamm N, Voipio-Pulkki LM, Pettersson K.](#)

[Related Articles, Links](#)

Molecular distinction of circulating pregnancy-associated plasma protein A in myocardial infarction and pregnancy.

Clin Chem. 2005 Jan;51(1):75-83.

PMID: 15613709 [PubMed - indexed for MEDLINE]

☐ 3: [Gerard N, Delpuech T, Oxvig C, Overgaard MT, Monget P.](#)

[Related Articles, Links](#)

Proteolytic degradation of IGF-binding protein (IGFBP)-2 in equine ovarian follicles: involvement of pregnancy-associated plasma protein-A (PAPP-A) and association with dominant but not subordinated follicles.

J Endocrinol. 2004 Sep;182(3):457-66.

PMID: 15350187 [PubMed - indexed for MEDLINE]

☐ 4: [Zorin NA, Zorina VN.](#)

[Related Articles, Links](#)

[Role of proteins of the macroglobulin family in the mechanisms of infection]

Zh Mikrobiol Epidemiol Immunobiol. 2004 May-Jun;(3):105-12.







Review. Russian.

PMID: 15346965 [PubMed - indexed for MEDLINE]

☐ 5: [Matsui M, Sonntag B, Hwang SS, Byerly T, Hourvitz A, Adashi EY, Shimasaki S.](#)


[Related Articles, Links](#)

Erickson GF.


-  Pregnancy-associated plasma protein-a production in rat granulosa cells: stimulation by follicle-stimulating hormone and inhibition by the oocyte-derived bone morphogenetic protein-15. *Endocrinology*. 2004 Aug;145(8):3686-95. Epub 2004 Apr 15. PMID: 15087430 [PubMed - indexed for MEDLINE]
- 6: Sivanandam AS, Mohan S, Kita H, Kapur S, Chen ST, Linkhart TA, Bagi G, Baylink DJ, Qin X. [Related Articles](#), [Links](#)
-  Studies on regulation of IGF (insulin-like growth factor)-binding protein (IGFBP) 4 proteolysis by pregnancy-associated plasma protein-A (PAPP-A) in cells treated with phorbol ester. *Biochem J*. 2004 Apr 1;379(Pt 1):57-64. PMID: 14705967 [PubMed - indexed for MEDLINE]
- 7: Christiansen M, Jaliashvili I. [Related Articles](#), [Links](#)
-  Total pregnancy-associated plasma protein A--a first trimester maternal serum marker for Down's syndrome: clinical and technical assessment of a poly-monoclonal enzyme immunoassay. *Scand J Clin Lab Invest*. 2003;63(6):407-15. PMID: 14594321 [PubMed - indexed for MEDLINE]
- 8: Arici A, Matalliotakis I, Goumenou A, Koumantakis G, Fragouli Y, Mahutte NG. [Related Articles](#), [Links](#)
-  Increased pregnancy-associated plasma protein-A (PAPP-A) concentrations in peritoneal fluid of women with endometriosis. *Am J Reprod Immunol*. 2003 Feb;49(2):70-4. PMID: 12765344 [PubMed - indexed for MEDLINE]
- 9: Rivera GM, Fortune JE. [Related Articles](#), [Links](#)
-  Selection of the dominant follicle and insulin-like growth factor (IGF)-binding proteins: evidence that pregnancy-associated plasma protein A contributes to proteolysis of IGF-binding protein 5 in bovine follicular fluid. *Endocrinology*. 2003 Feb;144(2):437-46. PMID: 12538602 [PubMed - indexed for MEDLINE]
- 10: Monget P, Mazerbourg S, Delpuech T, Maurel MC, Maniere S, Zapf J, Lalmanach G, Oxvig C, Overgaard MT. [Related Articles](#), [Links](#)
-  Pregnancy-associated plasma protein-A is involved in insulin-like growth factor binding protein-2 (IGFBP-2) proteolytic degradation in bovine and porcine preovulatory follicles: identification of

cleavage site and characterization of IGFBP-2 degradation.
Biol Reprod. 2003 Jan;68(1):77-86.
PMID: 12493698 [PubMed - indexed for MEDLINE]


- ☐ **11:** [Khosravi J, Diamandi A, Krishna RG, Bodani U, Mistry J, Khaja N.](#) Related Articles, Links

 Pregnancy associated plasma protein-A: ultrasensitive immunoassay and determination in coronary heart disease.
Clin Biochem. 2002 Oct;35(7):531-8.
PMID: 12493581 [PubMed - indexed for MEDLINE]


- ☐ **12:** [Sun IY, Overgaard MT, Oxvig C, Giudice LC.](#) Related Articles, Links

 Pregnancy-associated plasma protein A proteolytic activity is associated with the human placental trophoblast cell membrane.
J Clin Endocrinol Metab. 2002 Nov;87(11):5235-40.
PMID: 12414897 [PubMed - indexed for MEDLINE]


- ☐ **13:** [Qin X, Sexton C, Byun D, Strong DD, Baylink DJ, Mohan S.](#) Related Articles, Links

 Differential regulation of pregnancy associated plasma protein (PAPP)-A during pregnancy in human and mouse.
Growth Horm IGF Res. 2002 Oct;12(5):359-66.
PMID: 12213189 [PubMed - indexed for MEDLINE]


- ☐ **14:** [Qin QP, Christiansen M, Pettersson K.](#) Related Articles, Links

 Point-of-care time-resolved immunofluorometric assay for human pregnancy-associated plasma protein A: use in first-trimester screening for Down syndrome.
Clin Chem. 2002 Mar;48(3):473-83.
PMID: 11861437 [PubMed - indexed for MEDLINE]

- ☐ **15:** [Dumps P, Meisser A, Pons D, Morales MA, Anguenot JL, Campana A, Bischof P.](#) Related Articles, Links


 Accuracy of single measurements of pregnancy-associated plasma protein-A, human chorionic gonadotropin and progesterone in the diagnosis of early pregnancy failure.
Eur J Obstet Gynecol Reprod Biol. 2002 Jan 10;100(2):174-80.
PMID: 11750960 [PubMed - indexed for MEDLINE]

- ☐ **16:** [Byun D, Mohan S, Yoo M, Sexton C, Baylink DJ, Qin X.](#) Related Articles, Links


 Pregnancy-associated plasma protein-A accounts for the insulin-like growth factor (IGF)-binding protein-4 (IGFBP-4) proteolytic activity in human pregnancy serum and enhances the mitogenic

activity of IGF by degrading IGFBP-4 in vitro.
J Clin Endocrinol Metab. 2001 Feb;86(2):847-54.
PMID: 11158056 [PubMed - indexed for MEDLINE]


☐ **17:** [Qin X](#), [Byun D](#), [Lau KH](#), [Baylink DJ](#), [Mohan S](#). Related Articles, Links

 Evidence that the interaction between insulin-like growth factor (IGF)-II and IGF binding protein (IGFBP)-4 is essential for the action of the IGF-II-dependent IGFBP-4 protease.
Arch Biochem Biophys. 2000 Jul 15;379(2):209-16.
PMID: 10898936 [PubMed - indexed for MEDLINE]


☐ **18:** [Bersinger NA](#), [Meisser A](#), [Bessou T](#), [Seguin P](#), [Birkhauser MH](#), [Bischof P](#). Related Articles, Links

 Production and characterization of monoclonal antibodies against pregnancy-associated plasma protein A.
Mol Hum Reprod. 1999 Jul;5(7):675-81.
PMID: 10381824 [PubMed - indexed for MEDLINE]

☐ **19:** [Duan C](#), [Clemmons DR](#). Related Articles, Links

 Differential expression and biological effects of insulin-like growth factor-binding protein-4 and -5 in vascular smooth muscle cells.
J Biol Chem. 1998 Jul 3;273(27):16836-42.
PMID: 9642243 [PubMed - indexed for MEDLINE]

☐ **20:** [Rhoton-Vlasak A](#), [Wagner JM](#), [Rutgers JL](#), [Baergen RN](#), [Young RH](#), [Roche PC](#), [Plummer TB](#), [Gleich GJ](#). Related Articles, Links

 Placental site trophoblastic tumor: human placental lactogen and pregnancy-associated major basic protein as immunohistologic markers.
Hum Pathol. 1998 Mar;29(3):280-8.
PMID: 9496832 [PubMed - indexed for MEDLINE]

Items 1 - 20 of 49

Page of 3 Next

Display Show Sort by Send to


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 3 2005 04:38:48



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Display Show Sort by Send to

About Entrez

All: 49 Review: 2

Text Version

Items 21 - 40 of 49

Previous of 3 Next

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation
Matcher
Batch Citation
Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related
Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

- ☐ **21:** Qin QP, Christiansen M, Oxvig C, Pettersson K, Sottrup-Jensen L, Koch C, Norgaard-Pedersen B. [Related Articles, Links](#)



Double-monoclonal immunofluorometric assays for pregnancy-associated plasma protein A/proeosinophil major basic protein (PAPP-A/proMBP) complex in first-trimester maternal serum screening for Down syndrome.

Clin Chem. 1997 Dec;43(12):2323-32.

PMID: 9439450 [PubMed - indexed for MEDLINE]

- ☐ **22:** Qin Q, Christiansen M, Lovgren T, Norgaard-Pedersen B, Pettersson K. [Related Articles, Links](#)



Dual-label time-resolved immunofluorometric assay for simultaneous determination of pregnancy-associated plasma protein A and free beta-subunit of human chorionic gonadotrophin.

J Immunol Methods. 1997 Jul 14;205(2):169-75.

PMID: 9294599 [PubMed - indexed for MEDLINE]

- ☐ **23:** Qin QP, Nguyen TH, Christiansen M, Larsen SO, Norgaard-Pedersen B. [Related Articles, Links](#)



Time-resolved immunofluorometric assay of pregnancy-associated plasma protein A in maternal serum screening for Down's syndrome in first trimester of pregnancy.

Clin Chim Acta. 1996 Oct 29;254(2):113-29.

PMID: 8896900 [PubMed - indexed for MEDLINE]

- ☐ **24:** Casals E, Fortuny A, Grudzinskas JG, Suzuki Y, Teisner B, Comas C, Sanllehy C, Ojuel J, Borrell A, Soler A, Ballesta AM. [Related Articles, Links](#)



First-trimester biochemical screening for Down syndrome with the use of PAPP-A, AFP, and beta-hCG.

Prenat Diagn. 1996 May;16(5):405-10.

PMID: 8843997 [PubMed - indexed for MEDLINE]

- ❑ 25: [Bonno M, Oxvig C, Kephart GM, Wagner JM, Kristensen T, Sottrup-Jensen L, Gleich GJ](#) Related Articles, Links

Localization of pregnancy-associated plasma protein-A and colocalization of pregnancy-associated plasma protein-A messenger ribonucleic acid and eosinophil granule major basic protein messenger ribonucleic acid in placenta.

Lab Invest. 1994 Oct;71(4):560-6.

PMID: 7526035 [PubMed - indexed for MEDLINE]

- ❑ 26: [Bischof P, Meisser A](#) Related Articles, Links

Immunological heterogeneity of pregnancy-associated plasma protein-A (PAPP-A). Effects on the radioimmunoassay of PAPP-A. Br J Obstet Gynaecol. 1989 Jul;96(7):870-5.

PMID: 2475161 [PubMed - indexed for MEDLINE]

- ❑ 27: [Bueler MR, Bersinger NA](#) Related Articles, Links

Antiserum to pregnancy-associated plasma protein A (PAPP-A) recognizes human haptoglobin.

Br J Obstet Gynaecol. 1989 Jul;96(7):867-9.

PMID: 2475160 [PubMed - indexed for MEDLINE]

- ❑ 28: [Sinosich MJ, Saunders DM](#) Related Articles, Links

Potential role of pregnancy-associated plasma protein-A in human reproduction.

J Reprod Immunol. 1987 Jan;10(1):55-65.

PMID: 2438405 [PubMed - indexed for MEDLINE]

- ❑ 29: [Tornehave D, Chemnitz J, Westergaard JG, Teisner B, Poulsen HK, Bolton AE, Grudzinskas JG](#) Related Articles, Links

Placental proteins in peripheral blood and tissues of ectopic pregnancies.

Gynecol Obstet Invest. 1987;23(2):97-102.

PMID: 2438195 [PubMed - indexed for MEDLINE]

- ❑ 30: [Mowles EA, Pinto-Furtado LG, Bolton AE](#) Related Articles, Links

A two-site immunoradiometric assay for human pregnancy-associated plasma protein A (PAPP-A) using monoclonal antibodies.

J Immunol Methods. 1986 Dec 4;95(1):129-33.

PMID: 2431064 [PubMed - indexed for MEDLINE]

- ❑ **31:** [Chemnitz J, Folkersen J, Teisner B, Sinosich MJ, Tornehave D, Westergaard JG, Bolton AE, Grudzinskas JG.](#) [Related Articles, Links](#)



Comparison of different antibody preparations against pregnancy-associated plasma protein-A (PAPP-A) for use in localization and immunoassay studies.

Br J Obstet Gynaecol. 1986 Sep;93(9):916-23.

PMID: 2429686 [PubMed - indexed for MEDLINE]

- ❑ **32:** [Tornehave D, Folkersen J, Teisner B, Chemnitz J.](#) [Related Articles, Links](#)



Immunohistochemical aspects of immunological cross-reaction and masking of epitopes for localization studies on pregnancy-associated plasma protein A.

Histochem J. 1986 Apr;18(4):184-8.

PMID: 2426224 [PubMed - indexed for MEDLINE]

- ❑ **33:** [Udagawa Y, Armstrong SS, Waites GT, Bell SC, Horne CH, Thomson AW.](#) [Related Articles, Links](#)



Immunohistochemical localization of murine alpha 1-pregnancy-associated protein (alpha 1-PAP) in non-pregnant females: a comparative study with human pregnancy-associated alpha 2-glycoprotein (alpha 2-PAG).

Clin Exp Immunol. 1985 Aug;61(2):397-405.

PMID: 2412747 [PubMed - indexed for MEDLINE]

- ❑ **34:** [Sinosich MJ, Dodd J, Hudson CN, Tyler JR, Seppala M, Grudzinskas JG, Saunders DM.](#) [Related Articles, Links](#)



The influence of pergonal on in vitro production of placental protein 5 (PP5) by ovarian tumour cells.

Tumour Biol. 1985;6(3):233-42.

PMID: 2416032 [PubMed - indexed for MEDLINE]

- ❑ **35:** [Bersinger NA.](#) [Related Articles, Links](#)



Enzyme immunometric assay for the determination of pregnancy associated plasma protein A (PAPP-A) with the antigen as solid phase (conjoint IEMA).

Experientia. 1984 Sep 15;40(9):1022-4.

PMID: 6205894 [PubMed - indexed for MEDLINE]

- ❑ **36:** [Gore CH, Sutcliffe RG.](#) [Related Articles, Links](#)



Pregnancy-associated plasma protein A: purification under mild conditions, peptide mapping and tests for possible interactions with

trypsin, plasmin and complement.

Placenta. 1984 Jul-Aug;5(4):293-313.

PMID: 6209704 [PubMed - indexed for MEDLINE]

- ☐ **37:** [Dobashi K, Ajika K, Ohkawa T, Okano H, Okinaga S, Arai K.](#) Related Articles, Links



Immunohistochemical localization of pregnancy-associated plasma protein A (PAPP-A) in placentae from normal and pre-eclamptic pregnancies.

Placenta. 1984 May-Jun;5(3):205-12.

PMID: 6209702 [PubMed - indexed for MEDLINE]

- ☐ **38:** [Schindler AM, Bischof P.](#) Related Articles, Links



Histochemical localization of pregnancy-associated plasma protein A in fetal, infant, and adult organs and comparison between antisera.

Gynecol Obstet Invest. 1984;18(2):88-94.

PMID: 6207082 [PubMed - indexed for MEDLINE]

- ☐ **39:** [Bersinger NA.](#) Related Articles, Links



Two-site enzyme immunoassay for pregnancy associated plasma protein A (PAPP-A).

Clin Chim Acta. 1983 Nov 15;134(3):297-306.

PMID: 6196143 [PubMed - indexed for MEDLINE]

- ☐ **40:** [Pledger DR, Belfield A.](#) Related Articles, Links



The effect of excess antigen on the enzyme-linked immunosorbent assay of onco-placental proteins.

J Immunol Methods. 1983 Jul 15;61(2):227-32.

PMID: 6190948 [PubMed - indexed for MEDLINE]

Items 21 - 40 of 49

Previous 2 of 3 Next

Display Show Sort by Send to


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 3 2005 04:38:48



All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search for

Limits Preview/Index History Clipboard Details

Display Show Sort by Send to

About Entrez

All: 49 Review: 2

Text Version

Items 41 - 49 of 49

Previous 3 of 3

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation

Matcher

Batch Citation

Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

☐ **41:** Pledger DR, Belfield A.

Related Articles, Links



An ELISA for pregnancy-associated plasma protein A.
Ann Clin Biochem. 1983 Jul;20 (Pt 4):236-40.
PMID: 6193752 [PubMed - indexed for MEDLINE]

☐ **42:** Anthony F, Masson GM, Wood PJ.

Related Articles, Links



Development of a radioimmunoassay for pregnancy-associated plasma protein A and establishment of normal levels in the first trimester of pregnancy.
Ann Clin Biochem. 1983 Jan;20 Pt 1:26-30.
PMID: 6188409 [PubMed - indexed for MEDLINE]

☐ **43:** Sutcliffe RG, Kukulska-Langlands BM,

Related Articles, Links

Horne CH, Maclean AB, Jandial V,
Sutherland HW, Gibb S, Bowman AW.



Studies on the concentration of pregnancy-associated plasma protein-A during normal and complicated pregnancy.
Placenta. 1982 Jan-Mar;3(1):71-80.
PMID: 6176990 [PubMed - indexed for MEDLINE]

☐ **44:** Bischof P, Duberg S, Schindler AM.

Related Articles, Links



Is pregnancy-associated plasma protein A (PAPP-A) an immunomodulator during human pregnancy?
Placenta Suppl. 1982;4:93-102. No abstract available.
PMID: 6085897 [PubMed - indexed for MEDLINE]

☐ **45:** McIntyre JA, Hsi B, Faulk WP, Klopper A,

Related Articles, Links



Thomson R.
Immunological studies of the human placenta: functional and morphological analysis of pregnancy-associated plasma protein A (PAPP-A).
Immunology. 1981 Nov;44(3):577-83.

PMID: 6172371 [PubMed - indexed for MEDLINE]

- ☐ **46:** [Folkersen J, Grudzinskas JG, Hindersson P, Teisner B, Westergaard JG.](#) Related Articles, Links



Purification of pregnancy-associated plasma protein-A by a two step affinity chromatographic procedure.

Placenta. 1981 Jan-Mar;2(1):11-7.

PMID: 6163147 [PubMed - indexed for MEDLINE]

- ☐ **47:** [Sutcliffe RG, Kukulska-Langlands BM, Coggins JR, Hunter JB, Gore CH.](#) Related Articles, Links



Studies on human pregnancy-associated plasma protein A.

Purification by affinity chromatography and structural comparisons with alpha 2-macroglobulin.

Biochem J. 1980 Dec 1;191(3):799-809.

PMID: 6169340 [PubMed - indexed for MEDLINE]

- ☐ **48:** [Klopper A.](#) Related Articles, Links



Review article: the new placental proteins.

Placenta. 1980 Jan-Mar;1(1):77-89. Review. No abstract available.

PMID: 6160571 [PubMed - indexed for MEDLINE]

- ☐ **49:** [Bischof P, Bruce D, Cunningham P, Klopper A.](#) Related Articles, Links



Measurement of pregnancy associated plasma protein A (PAPP-A).

Clin Chim Acta. 1979 Jul 16;95(2):243-7.

PMID: 93522 [PubMed - indexed for MEDLINE]

Items 41 - 49 of 49

Previous **Page** 3 of 3

Display **Summary** Show **20** Sort by Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Nov 3 2005 04:38:48